Devartment of

Early childhood oral health needs

What is the Public Health Issue?

Oral health is integral to general health.¹ Early childhood caries (ECC), previously known as baby bottle tooth decay, is severe and rampant decay that occurs in the primary teeth of infants and toddlers. It affects 5-10% of America's young children,² and up to 20% of children

Consequences of untreated ECC: 2

- ✓ High risk of future decay⁵
- ✓ Pain
- ✓ Poor eating habits
- ✓ Weight loss
- ✓ Problems with speech, self-esteem, socialization

from low-income families.³ Less severe decay also occurs in young children. In fact, six out of 10 children have reported at least one or more decayed tooth by age five.⁴

ECC is usually associated with frequent overexposure of primary teeth to sugar. This often occurs when infants and toddlers are repeatedly given pacifying bottles or sippy cups filled with milk, juice or other sweet liquids during the day or at bedtime.¹ Another risk factor associated with ECC is the mother's oral health, and transmission of decay-causing bacteria from mother to child.²

The cost of treatment for ECC is very high. As of 2000, it was estimated that costs ranged from \$1,500 to \$2,000 per child depending on whether hospitalization and general anesthesia were necessary. Prevention of ECC requires a comprehensive plan involving "at risk" communities and parents, general and pediatric dental providers, and primary and pediatric non-dental providers. The emphasis should be on healthy feeding practices, daily tooth cleaning, appropriate use of toothpaste and other products that contain fluoride, and early identification and timely referral and intervention for at-risk children.

What is the situation in Maine?

In 2002, there were only ten dentists in Maine who specialized in pediatric dentistry. All of Maine's 16 counties include groups of communities designated as Dental Health Professional Shortage Areas.

Maine ECC Statistics, 1-4 year olds*

- ✓ 48% with tooth decay
- ✓ 13% with severe decay, ECC *from a survey of Washington County only

What is Maine doing?

• Maine's Early Childhood Caries Prevention and Intervention Program, currently under development, has the following three objectives: (1) increase oral health assessment of young pediatric patients by primary care providers (PCPs), (2) increase accessibility of oral health services for young pediatric patients,

Related Healthy People 2010 Objective:⁷

✓ Reduce tooth decay experience in 2-4 year olds to 11%.

Healthy Maine 2010 objective:8

✓ Reduce the proportion of children with tooth decay in their primary teeth to 25%

- and (3) increase appropriate dental referrals from PCPs to dental providers for follow-up services.
- Community water fluoridation prevents decay and provides a protective benefit to Maine's children. In 2004, 84% of the population in Maine on public water systems receives fluoridated water. However, this translates into only about 38.5% of Maine's total population receiving fluoridated water due to the large proportion of Maine citizens who drink water from a private source.

large proportion of Maine citizens who drink water from a private source.					

References:

- 1. U.S. Department of Health and Human Services. *Oral Health in America: a report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health 2000.
- 2. Amy Seif. Promoting Awareness, Preventing Pain: Facts on Early Childhood Caries. National Maternal and Child Oral Health Resource Center, 1999.
- 3. Bruerd B, Jones C, Krise D. Preventing baby bottle tooth decay and early childhood caries among AI/AN infants and children. The HIS Primary Care Provider 1997; 23 (3): 37-9.
- 4. U.S. Department of Health and Human Services, National Center for Health Statistics. National Health and Nutrition Examination Survey III, 1988-1994. CD-Rom Series 11, no. 1. Hyattsville, MD. National Center for Health Statistics, Centers for Disease Control and Prevention, 1997.
- 5. O'Sullivan DM, Tinanoff N. Maxillary anterior caries associated with increased risk in other primary teeth. Journal of Dental Research 1993; 72 (12): 1577-80.
- 6. Bureau of Health, Maine Department of Human Services. Maine Cooperative Health Manpower Resource System: General Practice and Specialist Dentists, 2002. http://www.maine.gov/dhs/bohodr/index.htm
- 7. US Department of Health and Human Services, Healthy People 2010. Washington DC: 2000. www.health.gov/healthypeople/
- 8. Bureau of Health, Maine Department of Human Services. Healthy Maine 2010: Longer and Healthier Lives. December 2002.

Published by the Oral Health Program, Bureau of Health, Maine Department of Health & Human Services, 2004, in cooperation with the Division of Oral Health, Centers for Disease Control and Prevention, Atlanta, Georgia. For more information, write to the OHP at 11 State House Station, Key Plaza, 4th floor, Augusta, ME 04333-0011, or call (207) 287-2361 or TTY (207) 287-8015 for the deaf or hard of hearing. Acknowledgements: Kathleen D. Heiden, RDH, MSPH, Barbara F. Gooch, DMD, MPH, Division of Oral Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, and Lisa Masinter, CDC ORISE Fellow, Summer 2003.



John Elias Baldacci, Governor John R. Nicholas, Commissioner